

Use of Amplidiag Bacterial GE assay (Mobidiag) in the diagnosis of bacterial diarrhoea

Saida Rasoanandrasana¹, Hervé Jacquier^{1,3}, Hana Benmansour¹, Philippe Bidet^{2,3}, Emmanuelle Cambau^{1,3}, Raskine Laurent¹, Stéphane Bonacorsi^{2,3}, Béatrice Berçot^{1,3}
 Laboratoire de bactériologie-virologie, Hôpital Lariboisière, Laboratoire de microbiologie, Hôpital Robert Debré, IAME, UMR 1137, INSERM, Université Paris Nord, Sorbonne Paris Cité, Paris, France


INTRODUCTION

The diagnosis of bacterial diarrhoea currently relies on cultures using selective media, a time-consuming approach whose sensitivity is still matter of debate, and time to result very long. Recently, several automated systems based on multiplex PCR assays have been developed to provide a rapid and comprehensive diagnosis. In this study, we assessed the Amplidiag Bacterial GE assay (Mobidiag) in the diagnosis of bacterial diarrhoea.

MATERIAL AND METHODS

100 consecutive stools collected between September and October 2015 in Lariboisière Hospital (Paris) for the diagnosis of infectious diarrhoea in adults, with a systematic screening of *Salmonella*, *Shigella* and *Campylobacter* by culture methods


43 frozen stools from children of Robert Debré Hospital (Paris), known to be positive for a large range of gastrointestinal pathogens covered by the kit



Bacterial pathogens detected by Amplidiag Bacterial GE assay (Mobidiag)

<i>Campylobacter</i>	EHEC
<i>Salmonella</i>	ETEC
Shigella/EIEC	EAEC
<i>Yersinia</i>	EPEC

RESULTS



43 frozen stools (RDB)

Identification obtained by culture	Detection of the cultured pathogen by PCR	Added microorganism detected by PCR
<i>Campylobacter</i> (n=12)	12	6
<i>Salmonella</i> (n=20)	15	10
<i>Shigella</i> (n=10)	10	5
<i>Yersinia</i> (n=1)	1	0

CONCLUSION

Amplidiag Bacterial GE assay is a **rapid and reliable method** to screen usual bacteria involved in gastroenteritis. This approach is particularly of interest to select samples that require cultures or not. Noteworthy, the high rate of pathogenic *E. coli* needs further investigations to unravel the clinical significance of *E. coli* positive results.